

袪朋官雉囫袞 B 橄蕃茑慈袪朋瞬倨禽橄蕃獭涌 鳞懒危釉除抹逡懒匝厦云潘情褰集

經吸鴉 鯪 铈 踞藏濱 贈 514071

[] B 豎

2012 6 誅2014 10 256 贈

斜128 斜128 豎 B

豎 姑 14姑28 d 豎

豎 斜 <0.05 豎 14姑28 d 贈

豎 66.4% 90.6%贈 39.8%姑74.3%斜 <0.05 豎 28 d 贈

豎 斜 <0.05 豎 B 豎

[] 閩喏鰈哇;標畹喏团襠;替得腴域涑懣 B 驟煤;替得飼愁岨驟煤

[] R758.2 [] A [] 1674-4721贈2015 10贈a -0133-03

Effect observion of the compound polymyxin B ointment combined with compound ointment flumetasone in treatment of chronic eczema and contact dermatitis

The Clinic in Huanan Road,Center for Disease Control and Prevention of Meizhou City in Guangdong Province,Meizhou 514071,China

[Abstract] Objective To observe the clinical effect of compound polymyxin B ointment combined with compound ointment flumetasone in treatment of chronic eczema and contact dermatitis. **Methods** 256 patients with chronic eczema and contact dermatitis in our center from June 2012 to October 2014 were selected, they were randomly divided into the study group (128 cases) and the control group (128 cases). The control group was treated with simple compound flumetasone ointment treatment, the study group was treated with compound ointment flumetasone and compound polymyxin B ointment treatment. The itching degree and other clinical symptoms score before and after treatment, clinical effect of after 14 and 28 d treatment, recurrence rate in two groups were observed. **Results** After treatment, the clinical symptoms in two groups were improved, the scores in study group were significantly lower than that of control group. After 14 and 28 d treatment, the Total effective rate in study group were respectively 66.4%, 90.6%, which were higher than 39.8% and 74.3% of control group. After 28 d treatment, the recurrence rate in study group was lower than that of control group, the differences were statistically significant (<0.05). **Conclusion** Compound polymyxin B ointment combined with compound ointment flumetasone combination therapy can significantly improve the symptoms of patients with chronic eczema and contact dermatitis, it has high effective rate, and it is suitable for long-term clinical application.

[Key words] Chronic eczema; Contact dermatitis; Compound polymyxin B ointment; Compound ointment flumetasone

贈 豎 B

[1]贈 豎

姑 豎 1 豎

[2-3]豎 1.1 癩籀铈

2012 6 誅2014 10 256 贈

贈 姑 贈 贈

噲 128 昱
噲 噲 噲
一 10%募 B姑
噲 昱 57 噲 71
噲 32誄60 噲 斜38.9^6.8 噲 1~5 噲
斜1.6^2.5 募 63 噲 65 噲 30誄66
噲 斜37.8^7.2 噲 2~6 噲 斜2.1^2.8 昱
姑 姑 噲
斜 >0.05 噲 昱

1.2 鐸陽

斜 噲 煖
HC20100009 噲 噲1 /d昱
B 斜
噲 煖H20061269 噲
B

1誄1 噲 噲1 /d昱 1
7d噲 2 昱

1.3 𪔐匀琨踝芳哖瞢程俞

姑 姑
斜 姑 4 姑 姑 姑 姑 姑 姑 姑 姑
姑 姑 4 0姑1姑2姑3 噲4
昱
[5]煖 / 2100%]
姑 姑 4 噲 姑 90%噲
61%~89%噲 咳20%姑
21%~60%噲

